

Table XXVIII-1

AGE-ADJUSTED SEER CANCER INCIDENCE^a AND U.S. DEATH^b RATES, 2000-2003

By Primary Cancer Site

All Races, Males and Females

Site	Ages 0-14						Ages 0-19					
	Incidence			Mortality			Incidence			Mortality		
	Total	Males	Females	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Sites												
All Races	14.8	15.7	13.9	2.5	2.7	2.3	16.4	17.4	15.2	2.8	3.1	2.5
Whites	15.7	16.7	14.7	2.6	2.7	2.4	17.5	18.8	16.2	2.8	3.1	2.5
Blacks	10.7	10.9	10.4	2.4	2.6	2.1	11.3	11.7	10.9	2.7	3.0	2.4
Bone & Joint	0.7	0.7	0.6	0.1	0.1	0.1	0.9	1.0	0.7	0.2	0.3	0.2
Brain & Other nervous	3.3	3.4	3.1	0.7	0.8	0.7	2.9	3.1	2.8	0.7	0.7	0.6
Hodgkin lymphoma	0.5	0.7	0.4	0.0	0.0		1.1	1.2	1.1	0.0	0.0	0.0
Kidney & Renal pelvis	0.8	0.8	0.8	0.1	0.1	0.1	0.6	0.6	0.6	0.1	0.1	0.1
Leukemia	4.8	5.1	4.5	0.8	0.9	0.7	4.3	4.7	3.9	0.9	1.0	0.8
Acute lymphocytic	3.8	4.0	3.5	0.4	0.4	0.3	3.2	3.5	2.9	0.4	0.5	0.3
Non-Hodgkin lymphoma	0.9	1.2	0.6	0.1	0.1	0.1	1.1	1.5	0.7	0.1	0.2	0.1
Soft tissue	1.0	1.0	1.0	0.1	0.1	0.1	1.1	1.1	1.0	0.2	0.2	0.2

^a SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^b Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Statistic could not be calculated. Rate based on less than 16 cases for the time interval.